
A HODAL APPROACH TO ARCTIC NOISE HODELLING HITH IMPLIED VERTICAL COHERENCE(U) NAVAL OCEAN SYSTEMS CENTER SAM DIEGO CA R B HILLIAMS FEB 88 AD-A192 966 1/1 UNCLASSIFIED F/G 20/2 NL



MICROPOPY RESOLUTION TEST CHART

		•	1
ntic	THE	СОРУ	
		LUPY	\leq

					<u>otic</u>	FILE	COPY U
Δ	D-A 19	2 066	PORT DOCUM	ENTATION PAGE			
~	D A 13	2 300		1b. RESTRICTIVE MARKIN	igs		
2a. SECURIT	Y CLASSIFICATION AL	JTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT			
2b. DECLASS	SIFICATION/DOWNGRA	ADING SCHEDULE		Approved for public r	elease; dístrib	ution is unlim	ited.
4. PERFORM	ING ORGANIZATION F	REPORT NUMBER (S	3)	5. MONITORING ORGANIZ	ATION REPORT	T NUMBER(S)	
			6b. OFFICE SYMBOL (# applicable) NOSC	7a. NAME OF MONITORING ORGANIZATION Naval Ocean Systems Center			
	an Systems Center S (City, State and ZIP Code)		14030	7b. ADDRESS (City, State and Z		.	
San Diego,	California 92152-5	5000		San Diego, California	a 92152-5000		
	F FUNDING/SPONSOR		(if applicable)	9. PROCUREMENT INSTR	UMENT IDENTIF	ICATION NUMB	ER
	an Research and De	velopment Activi	ty NORDA	10. 001/005 05 511/001/0			
BC. ADDRESS	S (City, State and ZIP Code)			10. SOURCE OF FUNDING PROGRAM ELEMENT NO.		TASK NO.	AGENCY ACCESSION NO.
NSTL, MS 39529-5004				62435N	MA63		DN307 376
11. TITLE (inc	lude Security Classification)						
A Modal	Approach to Arctic	Noise Modeling	with Implied Vertical	Coherence			
12. PERSON	AL AUTHOR(S)						
R.B. Will			· · · · · · · · · · · · · · · · · · ·				
13a. TYPE OF REPORT 13b. TIME COVERE			14. DATE OF REPORT (Year, Month, Day) 15. PAGE COUNT		NT		
Presentation 16. SUPPLEM	on/speech	FROM Nov 1987	7 TO Nov 1987	February 1988			
							;
17. COSATI C			18. SUBJECT TERMS (Continue on reverse if necessary and idea	ntifiy by block number)		
FIELD	GROUP	SUB-GROUP	modal energy wavelength correlation lengths				
		_					
							
19. ABSTRAC	CT (Continue on reverse if nece	essary and identify by block n	umber)				
tioning of a bution func 50 Hz have ally has a s two waveler although of noise will c sensitivity c	mbient noise source tions to calculate va- been analyzed for in x/x dependence angths are observed. her interpretations a ause the correlation calculations, however	es are proposed. In the state of the state o	These functions can be umbient noise function not using a bandwidth ro crossing at one was in lengths are consisted at simple model calcule several more distance fur e model validity. [Wo	ate for deep Arctic under be used, together with va- ns, such as vertical cohe of about 5 Hz. These di- velength. Sometimes, ho- ent with much of the noi- ulations show agreement at sources support the on- nction by itself is not a goork supported by NORDA	rious assumed rence. Vertica at a show that wowever, zero c see energy radi with the data. He wavelength good validator A and NOSC (I spatial and to il noise data be the vertical co- rossing lengths ating from a s A strong nea zero crossing to of the model to 6.2 Arctic pro	emporal distri- etween 15 and herence usu- s of more than mall region, rby source of result. Initial functions. grams.]
Presented	at the Acoustical Sc	ociety of America	, 16-20 November 19				TC ECTE 25 1988

20. DISTRIBUTION/AVAILABILITY OF ABSTRACT 21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED/UNLIMITED X SAME AS RPT DTIC USERS UNCLASSIFIED 22a. NAME OF RESPONSIBLE INDIVIDUAL 22b. TELEPHONE (include Area Code) 22c. OFFICE SYMBOL R.B. Williams (619) 553-3096 Code 541

M4. A modal approach to Arctic noise modeling with implied vertical coherence. R. B. Williams (Naval Ocean Systems Center, San Diego, CA 92152)

Analytic formulas of the modal attenuation coefficients appropriate for deep Arctic under-ice conditions, and modal energy partitioning of ambient noise sources are proposed. These functions can be used, together with various assumed spatial and temporal distribution functions to calculate various statistical ambient noise functions, such as vertical coherence. Vertical noise data between 15 and 50 Hz have been analyzed for model comparison, using a bandwidth of about 5 Hz. These data show that the vertical coherence usually has a $\sin x/x$ dependence, with the first zero crossing at one wavelength. Sometimes, however, zero crossing lengths of more than two wavelengths are observed. Longer correlation lengths are consistent with much of the noise energy radiating from a small region, although other interpretations are possible. Initial simple model calculations show agreement with the data. A strong nearby source of noise will cause the correlation to lengthen, while several more distant sources support the one wavelength zero crossing result. Initial sensitivity calculations, however, show that the vertical coherence function by itself is not a good validator of the model functions. Other statistics will be sought to further examine model validity. [Work supported by NORDA and NOSC 6.2 Arctic programs.]

Acces	sion For	
DTIC	GRARI TAB counced fication	
By	·ibutien/	
Ava	lability	Codes
Dist	Aveil es Specie	•
A-1		

END DATE FILMED DT/C July 88